Jefferson County

Youth & Tobacco Data Summary 10th Grade

Year 2000

10th Grade Students Surveyed n= 186			Year 2000	County %	State %
Female	10th Grade S	Students Surveyed	n= 186	100.0 %	100.0 %
Male 51.9 % 48.4 % Race/Ethnicity Students of Color White 90.2 % 75.8 % County %+/- Margin of Error State %+/- Margin of Error Past 30 days, youth who have smoked cigarettes at least once Female n= 89 39.3 %+/- 10.1 21.8 %+/- 2.2 Male n= 96 14.6 %+/- 7.1 17.5 %+/- 2.1 Total n= 185 26.3 %+/- 6.3 19.8 %+/- 1.5 Past 30 days, youth who have smoked cigars at least once Emale n= 89 2.2 %+/- 3.1 4.8 %+/- 1.2 Male n= 95 11.6 %+/- 6.4 11.2 %+/- 1.7 Total n= 184 0.0 %+/- 0.0 1.2 %+/- 0.6 Past 30 days, youth who have smoked bidis (flavored cigarettes) at least once Female n= 86 0.0 %+/- 0.0 1.2 %+/- 0.5 Past 30 days, youth who have used smokeless tobacco (chewing tobacco, snuff or dip) at least once Female n= 86 2.3 %+/- 3.1 4.5 %+/- 1.2 Male n= 96 1.1 %+/- 2.2 1.4 %+/- 0.8 <	Gender				
Race/Ethnicity Students of Color White 90.2 % 75.8 % 24.2 % 75.8 % 25.0 % 75.8 % 7	Female			48.1 %	51.6 %
Students of Color 9.8 % 94.2 % 75.8 % 90.2 % 75.8 % 75.8 % 90.2 % 75.8 % 75.8 % 90.2 % 75.8 %	Male			51.9 %	48.4 %
White 90.2 % 75.8 % County %+/- Margin of Error State %+/- Margin of Error Past 30 days, youth who have smoked cigarettes at least once 39.3 %+/- 10.1 21.8 %+/- 2.2 Female n = 89 39.3 %+/- 10.1 21.8 %+/- 2.1 Total n = 185 26.3 %+/- 6.3 19.8 %+/- 1.5 Past 30 days, youth who have smoked cigars at least once Female n = 89 2.2 %+/- 3.1 4.8 %+/- 1.2 Female n = 89 2.2 %+/- 3.1 4.8 %+/- 1.7 1.7 7.0 %+/- 3.7 7.9 %+/- 1.7 Total n = 184 7.0 %+/- 3.7 7.9 %+/- 1.0 7.9 %+/- 1.0 Past 30 days, youth who have smoked tobacco in a pipe at least once 8 0.0 %+/- 0.0 1.2 %+/- 0.6 Female n = 88 0.0 %+/- 0.0 1.2 %+/- 0.8 2.7 %+/- 0.8 Past 30 days, youth who have smoked bidis (flavored cigarettes) at least once 2 2.3 %+/- 1.1 4.5 %+/- 1.1 Female n = 88 2.3 %+/- 3.1 4.5 %+/- 1.5 2.7 %+/- 0.8 Past 30 days, youth who have used smokeless tobacco (chewing tobacco, snuff or dip) at least once <t< td=""><td>Race/Ethnici</td><td>ty</td><td></td><td></td><td></td></t<>	Race/Ethnici	ty			
County %+/- Margin of Error	Students of	Color		9.8 %	24.2 %
Margin of Error Margin of Error Margin of Error	White			90.2 %	75.8 %
Margin of Error Margin of Error Margin of Error				County %+/-	State %+/-
Past 30 days, youth who have smoked cigarettes at least once Female n = 89 39.3 %+/- 10.1 21.8 %+/- 2.2 Male n = 96 14.6 %+/- 7.1 17.5 %+/- 2.1 Total n = 185 26.3 %+/- 6.3 19.8 %+/- 1.5 Past 30 days, youth who have smoked cigars at least once Female n = 89 2.2 %+/- 3.1 4.8 %+/- 1.2 Male n = 95 11.6 %+/- 6.4 11.2 %+/- 1.7 Total n = 184 7.0 %+/- 3.7 7.9 %+/- 1.0 Past 30 days, youth who have smoked tobacco in a pipe at least once Female n = 88 0.0 %+/- 0.0 1.2 %+/- 0.6 Male n = 96 3.1 %+/- 3.5 2.7 %+/- 0.5 Past 30 days, youth who have smoked bidis (flavored cigarettes) at least once Female n = 88 2.3 %+/- 1.8 1.9 %+/- 0.5 Past 30 days, youth who have smoked bidis (flavored cigarettes) at least once Female n = 88 2.3 %+/- 3.1 4.5 %+/- 1.1 Male n = 96 5.2 %+/- 4.4 4.7 %+/- 1.2 Total n = 184 3.8 %+/- 2.7 4.6 %+/- 0.8 Past 30 days, youth who have used smokeless tobacco (chewing tobacco, snuff or dip) at least once Female n = 87 1.1 %+/- 2.2 1.4 %+/- 0.8 Past 30 days, youth who have used smokeless tobacco (chewing tobacco, snuff or dip) at least once Female n = 87 1.1 %+/- 2.2 1.4 %+/- 0.8 Past 30 days, youth who have used any tobacco product at least once Female n = 183 6.5 %+/- 3.6 4.6 %+/- 0.9 Past 30 days, youth who have used any tobacco product at least once Female n = 88 42.0 %+/- 10.3 23.8 %+/- 2.3 44.8 %+/- 2.4 4					Margin
Past 30 days, youth who have smoked cigarettes at least once Female n = 89 Male n = 96 Total n = 185 Past 30 days, youth who have smoked cigars at least once Female n = 89 Ale n = 96 Female n = 89 Ale n = 96 Ale n = 96 Ale n = 96 Ale n = 89 Ale n = 89 Ale n = 89 Ale n = 95 Ale n = 184 Past 30 days, youth who have smoked cigars at least once Female n = 184 Past 30 days, youth who have smoked tobacco in a pipe at least once Female n = 88 Ale n = 96 Ale n = 96 Ale n = 184 Past 30 days, youth who have smoked tobacco in a pipe at least once Female n = 88 Ale n = 96 Ale n = 96 Ale n = 184 Past 30 days, youth who have smoked bidis (flavored cigarettes) at least once Female n = 184 Past 30 days, youth who have smoked bidis (flavored cigarettes) at least once Female n = 88 Ale n = 96 Ale n = 184 Past 30 days, youth who have used smokeless tobacco (chewing tobacco, snuff or dip) at least once Female n = 87 Ale n = 96 Ale n =				_	_
Female			oked cigarettes		5. —
Male n= 96 14.6 %+/- 7.1 17.5 %+/- 2.1 17.5 %+/- 2.1 15.5 %+/- 2.1 18.8 %+/- 1.5 19.8 %+/- 1.5 Past 30 days, youth who have smoked cigars at least once Female n= 89 2.2 %+/- 3.1 4.8 %+/- 1.2 4.7 Total 11.6 %+/- 6.4 11.2 %+/- 1.7 1.7 Total n= 184 7.0 %+/- 3.7 7.9 %+/- 1.0 Past 30 days, youth who have smoked tobacco in a pipe at least once n= 88 0.0 %+/- 0.0 1.2 %+/- 0.6 1.2 %+/- 0.8				39.3 %+/- 10.1	21.8 %+/- 2.2
Total n= 185 26.3 %+/- 6.3 19.8 %+/- 1.5 Past 30 days, youth who have smoked cigars at least once Female n= 89 2.2 %+/- 3.1 4.8 %+/- 1.2 Male n= 95 11.6 %+/- 6.4 11.2 %+/- 1.7 Total n= 184 7.0 %+/- 3.7 7.9 %+/- 1.0 Past 30 days, youth who have smoked tobacco in a pipe at least once Female n= 88 0.0 %+/- 0.0 1.2 %+/- 0.6 Male n= 96 3.1 %+/- 3.5 2.7 %+/- 0.8 Total n= 184 1.6 %+/- 1.8 1.9 %+/- 0.5 Past 30 days, youth who have smoked bidis (flavored cigarettes) at least once Female n= 88 2.3 %+/- 3.1 4.5 %+/- 0.5 Past 30 days, youth who have used smokeless tobacco (chewing tobacco, snuff or dip) at least once Female n= 87 1.1 %+/- 2.2 1.4 %+/- 0.8 Male n= 96 11.5 %+/- 6.4 7.9 %+/- 10. Total n= 183 6.5 %+/- 3.6 4.6 %+/- 0.9 Past 30 days, youth who have used any tobacco product at least once	Male	n= 96		14.6 %+/- 7.1	17.5 %+/- 2.1
Female n = 89 2.2 %+/- 3.1 4.8 %+/- 1.2 Male n = 95 11.6 %+/- 6.4 11.2 %+/- 1.7 Total n = 184 7.0 %+/- 3.7 7.9 %+/- 1.0 Past 30 days, youth who have smoked tobacco in a pipe at least once Female n = 88 0.0 %+/- 0.0 1.2 %+/- 0.6 Male n = 96 3.1 %+/- 3.5 2.7 %+/- 0.8 Total n = 184 1.6 %+/- 1.8 1.9 %+/- 0.5 Past 30 days, youth who have smoked bidis (flavored cigarettes) at least once Female n = 88 2.3 %+/- 3.1 4.5 %+/- 1.1 Male n = 96 5.2 %+/- 4.4 4.7 %+/- 1.2 Total n = 184 3.8 %+/- 2.7 4.6 %+/- 0.8 Past 30 days, youth who have used smokeless tobacco (chewing tobacco, snuff or dip) at least once Female n = 87 1.1 %+/- 2.2 1.4 %+/- 0.8 Male n = 96 11.5 %+/- 6.4 7.9 %+/- 1.6 Total n = 183 6.5 %+/- 3.6 4.6 %+/- 0.9 Past 30 days, youth who have used any tobacco product at least once Female n = 88 42.0 %+/- 10.3 23.8 %+/- 2.3	Total	n= 185			
Female n = 89 2.2 %+/- 3.1 4.8 %+/- 1.2 Male n = 95 11.6 %+/- 6.4 11.2 %+/- 1.7 Total n = 184 7.0 %+/- 3.7 7.9 %+/- 1.0 Past 30 days, youth who have smoked tobacco in a pipe at least once Female n = 88 0.0 %+/- 0.0 1.2 %+/- 0.6 Male n = 96 3.1 %+/- 3.5 2.7 %+/- 0.8 Total n = 184 1.6 %+/- 1.8 1.9 %+/- 0.5 Past 30 days, youth who have smoked bidis (flavored cigarettes) at least once Female n = 88 2.3 %+/- 3.1 4.5 %+/- 1.1 Male n = 96 5.2 %+/- 4.4 4.7 %+/- 1.2 Total n = 184 3.8 %+/- 2.7 4.6 %+/- 0.8 Past 30 days, youth who have used smokeless tobacco (chewing tobacco, snuff or dip) at least once Female n = 87 1.1 %+/- 2.2 1.4 %+/- 0.8 Male n = 96 11.5 %+/- 6.4 7.9 %+/- 1.6 Total n = 183 6.5 %+/- 3.6 4.6 %+/- 0.9 Past 30 days, youth who have used any tobacco product at least once Female n = 88 42.0 %+/- 10.3 23.8 %+/- 2.3	Past 30 days	, youth who have smo	oked cigars at least once		
Male n= 95 11.6 %+/- 6.4 11.2 %+/- 1.7 1.0 Total n= 184 7.0 %+/- 3.7 7.9 %+/- 1.0 Past 30 days, youth who have smoked tobacco in a pipe at least once Female n= 88 0.0 %+/- 0.0 1.2 %+/- 0.6 Male n= 96 3.1 %+/- 3.5 2.7 %+/- 0.8 Total n= 184 1.6 %+/- 1.8 1.9 %+/- 0.5 Past 30 days, youth who have smoked bidis (flavored cigarettes) at least once Female n= 88 2.3 %+/- 3.1 4.5 %+/- 1.1 Male n= 96 5.2 %+/- 4.4 4.7 %+/- 1.2 Total n= 184 3.8 %+/- 2.7 4.6 %+/- 0.8 Past 30 days, youth who have used smokeless tobacco (chewing tobacco, snuff or dip) at least once Female n= 87 1.1 %+/- 2.2 1.4 %+/- 0.8 Male n= 96 11.5 %+/- 6.4 7.9 %+/- 10.6 Total n= 183 6.5 %+/- 3.6 4.6 %+/- 0.9 Past 30 days, youth who have used any tobacco product at least once Female n= 88 42.0 %+/- 10.3 23.8 %+/- 2.3 Male n= 96 25.0 %+/- 8.7	•	· •	3	2.2 %+/- 3.1	4.8 %+/- 1.2
Total n= 184 7.0 %+/- 3.7 7.9 %+/- 1.0 Past 30 days, youth who have smoked tobacco in a pipe at least once Female n= 88 Male n= 96 Total n= 184 0.0 %+/- 0.0 1.2 %+/- 0.6 3.1 %+/- 3.5 2.7 %+/- 0.8 Total n= 184 1.6 %+/- 1.8 1.9 %+/- 0.5 Past 30 days, youth who have smoked bidis (flavored cigarettes) at least once Female n= 88 Male n= 96 2.3 %+/- 3.1 4.5 %+/- 1.1 1.0 %+/- 1.2 1.1 %+/- 2.7 1.1 %+/- 2.7 1.1 %+/- 2.2 1.4 %+/- 0.8 Past 30 days, youth who have used smokeless tobacco (chewing tobacco, snuff or dip) at least once Female n= 87 Male n= 96 1.1 %+/- 2.2 1.4 %+/- 0.8 Past 30 days, youth who have used any tobacco product at least once Female n= 183 Past 30 days, youth who have used any tobacco product at least once Female n= 88 Male n= 96 42.0 %+/- 10.3 23.8 %+/- 2.3 Male n= 96					
at least once Female n= 88 0.0 %+/- 0.0 1.2 %+/- 0.6 Male n= 96 3.1 %+/- 3.5 2.7 %+/- 0.8 Total n= 184 1.6 %+/- 1.8 1.9 %+/- 0.5 Past 30 days, youth who have smoked bidis (flavored cigarettes) at least once Female n= 88 2.3 %+/- 3.1 4.5 %+/- 1.1 Male n= 96 5.2 %+/- 4.4 4.7 %+/- 1.2 Total n= 184 3.8 %+/- 2.7 4.6 %+/- 0.8 Past 30 days, youth who have used smokeless tobacco (chewing tobacco, snuff or dip) at least once Female n= 87 1.1 %+/- 2.2 1.4 %+/- 0.8 Male n= 96 11.5 %+/- 6.4 7.9 %+/- 1.6 Total n= 183 6.5 %+/- 3.6 4.6 %+/- 0.9 Past 30 days, youth who have used any tobacco product at least once Female n= 88 42.0 %+/- 10.3 23.8 %+/- 2.3 Male n= 96 42.0 %+/- 10.3 23.8 %+/- 2.3 Male n= 96 25.0 %+/- 8.7 24.8 %+/- 2.4					
at least once Female n= 88 0.0 %+/- 0.0 1.2 %+/- 0.6 Male n= 96 3.1 %+/- 3.5 2.7 %+/- 0.8 Total n= 184 1.6 %+/- 1.8 1.9 %+/- 0.5 Past 30 days, youth who have smoked bidis (flavored cigarettes) at least once Female n= 88 2.3 %+/- 3.1 4.5 %+/- 1.1 Male n= 96 5.2 %+/- 4.4 4.7 %+/- 1.2 Total n= 184 3.8 %+/- 2.7 4.6 %+/- 0.8 Past 30 days, youth who have used smokeless tobacco (chewing tobacco, snuff or dip) at least once Female n= 87 1.1 %+/- 2.2 1.4 %+/- 0.8 Male n= 96 11.5 %+/- 6.4 7.9 %+/- 1.6 Total n= 183 6.5 %+/- 3.6 4.6 %+/- 0.9 Past 30 days, youth who have used any tobacco product at least once Female n= 88 42.0 %+/- 10.3 23.8 %+/- 2.3 Male n= 96 42.0 %+/- 10.3 23.8 %+/- 2.3 Male n= 96 25.0 %+/- 8.7 24.8 %+/- 2.4	Past 30 days	, youth who have smo	oked tobacco in a pipe		
Male Total n= 96 3.1 %+/- 3.5 2.7 %+/- 0.8 Past 30 days, youth who have smoked bidis (flavored cigarettes) at least once Female n= 88 2.3 %+/- 3.1 4.5 %+/- 1.1 Male n= 96 5.2 %+/- 4.4 4.7 %+/- 1.2 Total n= 184 3.8 %+/- 2.7 4.6 %+/- 0.8 Past 30 days, youth who have used smokeless tobacco (chewing tobacco, snuff or dip) at least once Female n= 87 1.1 %+/- 2.2 1.4 %+/- 0.8 Male n= 96 11.5 %+/- 6.4 7.9 %+/- 1.6 Total n= 183 6.5 %+/- 3.6 4.6 %+/- 0.9 Past 30 days, youth who have used any tobacco product at least once Female n= 88 42.0 %+/- 10.3 23.8 %+/- 2.3 Male n= 96 25.0 %+/- 8.7 24.8 %+/- 2.4	-		• •		
Male Total n= 96 3.1 %+/- 3.5 2.7 %+/- 0.8 Past 30 days, youth who have smoked bidis (flavored cigarettes) at least once Female n= 88 2.3 %+/- 3.1 4.5 %+/- 1.1 Male n= 96 5.2 %+/- 4.4 4.7 %+/- 1.2 Total n= 184 3.8 %+/- 2.7 4.6 %+/- 0.8 Past 30 days, youth who have used smokeless tobacco (chewing tobacco, snuff or dip) at least once Female n= 87 1.1 %+/- 2.2 1.4 %+/- 0.8 Male n= 96 11.5 %+/- 6.4 7.9 %+/- 1.6 Total n= 183 6.5 %+/- 3.6 4.6 %+/- 0.9 Past 30 days, youth who have used any tobacco product at least once Female n= 88 42.0 %+/- 10.3 23.8 %+/- 2.3 Male n= 96 25.0 %+/- 8.7 24.8 %+/- 2.4	Female	n= 88		0.0 %+/- 0.0	1.2 %+/- 0.6
Total 1.6 %+/- 1.8 1.9 %+/- 0.5 Past 30 days, youth who have smoked bidis (flavored cigarettes) at least once Female n= 88 2.3 %+/- 3.1 4.5 %+/- 1.1 Male n= 96 5.2 %+/- 4.4 4.7 %+/- 1.2 Total n= 184 3.8 %+/- 2.7 4.6 %+/- 0.8 Past 30 days, youth who have used smokeless tobacco (chewing tobacco, snuff or dip) at least once Female n= 87 1.1 %+/- 2.2 1.4 %+/- 0.8 Male n= 96 11.5 %+/- 6.4 7.9 %+/- 1.6 Total n= 183 6.5 %+/- 3.6 4.6 %+/- 0.9 Past 30 days, youth who have used any tobacco product at least once Female n= 88 42.0 %+/- 10.3 23.8 %+/- 2.3 Male n= 96 25.0 %+/- 8.7 24.8 %+/- 2.4		n= 96			
at least once Female n= 88 2.3 %+/- 3.1 4.5 %+/- 1.1 Male n= 96 5.2 %+/- 4.4 4.7 %+/- 1.2 Total n= 184 3.8 %+/- 2.7 4.6 %+/- 0.8 Past 30 days, youth who have used smokeless tobacco (chewing tobacco, snuff or dip) at least once Female n= 87 1.1 %+/- 2.2 1.4 %+/- 0.8 Male n= 96 11.5 %+/- 6.4 7.9 %+/- 1.6 Total n= 183 6.5 %+/- 3.6 4.6 %+/- 0.9 Past 30 days, youth who have used any tobacco product at least once Female n= 88 42.0 %+/- 10.3 23.8 %+/- 2.3 Male n= 96 25.0 %+/- 8.7 24.8 %+/- 2.4					
at least once Female n= 88 2.3 %+/- 3.1 4.5 %+/- 1.1 Male n= 96 5.2 %+/- 4.4 4.7 %+/- 1.2 Total n= 184 3.8 %+/- 2.7 4.6 %+/- 0.8 Past 30 days, youth who have used smokeless tobacco (chewing tobacco, snuff or dip) at least once Female n= 87 1.1 %+/- 2.2 1.4 %+/- 0.8 Male n= 96 11.5 %+/- 6.4 7.9 %+/- 1.6 Total n= 183 6.5 %+/- 3.6 4.6 %+/- 0.9 Past 30 days, youth who have used any tobacco product at least once Female n= 88 42.0 %+/- 10.3 23.8 %+/- 2.3 Male n= 96 25.0 %+/- 8.7 24.8 %+/- 2.4	Past 30 days	, youth who have smo	oked bidis (flavored cigarettes)		
Male Total n= 96 n= 184 5.2 %+/- 4.4 4.7 %+/- 1.2 Past 30 days, youth who have used smokeless tobacco (chewing tobacco, snuff or dip) at least once Female n= 87 n= 87 n= 96 n= 11.5 %+/- 6.4 n= 183 1.1 %+/- 2.2 n= 1.4 %+/- 0.8 n= 1.6 n= 183 Past 30 days, youth who have used any tobacco product at least once 6.5 %+/- 3.6 n= 10.3 n	-	• •	,		
Total 3.8 %+/- 2.7 4.6 %+/- 0.8 Past 30 days, youth who have used smokeless tobacco (chewing tobacco, snuff or dip) at least once Female n= 87 1.1 %+/- 2.2 1.4 %+/- 0.8 Male n= 96 11.5 %+/- 6.4 7.9 %+/- 1.6 Total n= 183 6.5 %+/- 3.6 4.6 %+/- 0.9 Past 30 days, youth who have used any tobacco product at least once Female n= 88 42.0 %+/- 10.3 23.8 %+/- 2.3 Male n= 96 25.0 %+/- 8.7 24.8 %+/- 2.4	Female	n= 88		2.3 %+/- 3.1	4.5 %+/- 1.1
Total 3.8 %+/- 2.7 4.6 %+/- 0.8 Past 30 days, youth who have used smokeless tobacco (chewing tobacco, snuff or dip) at least once Female n= 87 1.1 %+/- 2.2 1.4 %+/- 0.8 Male n= 96 11.5 %+/- 6.4 7.9 %+/- 1.6 Total n= 183 6.5 %+/- 3.6 4.6 %+/- 0.9 Past 30 days, youth who have used any tobacco product at least once Female n= 88 42.0 %+/- 10.3 23.8 %+/- 2.3 Male n= 96 25.0 %+/- 8.7 24.8 %+/- 2.4	Male	n= 96		5.2 %+/- 4.4	4.7 %+/- 1.2
(chewing tobacco, snuff or dip) at least once Female n= 87 1.1 %+/- 2.2 1.4 %+/- 0.8 Male n= 96 11.5 %+/- 6.4 7.9 %+/- 1.6 Total n= 183 6.5 %+/- 3.6 4.6 %+/- 0.9 Past 30 days, youth who have used any tobacco product at least once Female n= 88 42.0 %+/- 10.3 23.8 %+/- 2.3 Male n= 96 25.0 %+/- 8.7 24.8 %+/- 2.4					
(chewing tobacco, snuff or dip) at least once Female n= 87 1.1 %+/- 2.2 1.4 %+/- 0.8 Male n= 96 11.5 %+/- 6.4 7.9 %+/- 1.6 Total n= 183 6.5 %+/- 3.6 4.6 %+/- 0.9 Past 30 days, youth who have used any tobacco product at least once Female n= 88 42.0 %+/- 10.3 23.8 %+/- 2.3 Male n= 96 25.0 %+/- 8.7 24.8 %+/- 2.4	Past 30 days	, youth who have use	d smokeless tobacco		
Female n= 87 1.1 %+/- 2.2 1.4 %+/- 0.8 Male n= 96 11.5 %+/- 6.4 7.9 %+/- 1.6 Total n= 183 6.5 %+/- 3.6 4.6 %+/- 0.9 Past 30 days, youth who have used any tobacco product at least once Female n= 88 42.0 %+/- 10.3 23.8 %+/- 2.3 Male n= 96 25.0 %+/- 8.7 24.8 %+/- 2.4					
Male Total n= 96 n= 183 11.5 %+/- 6.4 7.9 %+/- 1.6 Past 30 days, youth who have used any tobacco product at least once 42.0 %+/- 10.3 23.8 %+/- 2.3 Female Nale n= 96 42.0 %+/- 8.7 24.8 %+/- 2.4	•	• • •		1.1 %+/- 2.2	1.4 %+/- 0.8
Total n= 183 6.5 %+/- 3.6 4.6 %+/- 0.9 Past 30 days, youth who have used any tobacco product at least once Female n= 88 42.0 %+/- 10.3 23.8 %+/- 2.3 Male n= 96 25.0 %+/- 8.7 24.8 %+/- 2.4	Male	n= 96		11.5 %+/- 6.4	
at least once Female n= 88 Male n= 96 42.0 %+/- 10.3 23.8 %+/- 2.3 25.0 %+/- 8.7 24.8 %+/- 2.4					
Female n= 88 42.0 %+/- 10.3 23.8 %+/- 2.3 Male n= 96 25.0 %+/- 8.7 24.8 %+/- 2.4	Past 30 days	, youth who have use	d any tobacco product		
Male n= 96 <u>25.0 %+/- 8.7</u> 24.8 %+/- 2.4	at least onc	e			
	Female	n= 88		42.0 %+/- 10.3	23.8 %+/- 2.3
	Male	n= 96		25.0 %+/- 8.7	24.8 %+/- 2.4
	Total	n= 184		33.0 %+/- 6.8	24.4 %+/- 1.6

Source: Washington State Department of Health Tobacco Prevention and Control

Assessment and Evaluation

n = number of survey respondents for county.

* = Rate not calculated because number of responses was less than 15.

Jefferson County

Youth & Tobacco Data Summary 10th Grade Year 2000

	County %+/- Margin of Error	State %+/- Margin of Error
About how many cigarettes have you smoked in your entire life? None 1/2 pack or less 1 - 4 packs 5 or more packs	n= 180 44.4 %+/- 7.3 22.8 %+/- 6.1 13.3 %+/- 5.0 19.4 %+/- 5.8	49.9 %+/- 1.9 24.7 %+/- 1.6 9.3 %+/- 1.1 16.1 %+/- 1.4
Do you want to stop using tobacco right now? (among current tobacco users) Yes	n= 52 63.5 %+/- 13.1	56.8 %+/- 4.0
Have you ever participated in a program to help you quit using tobacco? (among current tobacco users) Yes	n= 65 7.7 %+/- 6.5	6.7 %+/- 1.7
During past 12 months, did you ever try to quit using tobacco? (cigarettes, cigars, or chew/dip)(among current tobacco users) Yes	n= 33 57.6 %+/- 16.9	58.3 %+/- 4.9
Do you think you will smoke a cigarette at any time during the next year? Definitely no	n= 185 51.9 %+/- 7.2	56.2 %+/- 1.8
If one of your best friends offered you a cigarette, would you smoke it? Definitely no	n= 185 56.2 %+/- 7.1	59.1 %+/- 1.8
Susceptibility Index (combination of the previous two questions, either of which receives any answer other than 'Definitely no') Susceptible	n= 185 50.8 %+/- 7.2	46.8 %+/- 1.8
If one of your best friends offered you a cigarette, would you smoke it? (from among youth who did not smoke in last 30 days) Definitely no	n= 136 74.3 %+/- 7.3	72.8 %+/- 1.8
Do you think the smoke from other people's cigarettes (second-hand smoke) is harmful to you? Definitely yes or Probably yes	n= 185 92.4 %+/- 3.8	92.2 %+/- 1.0
Do you think it is safe to smoke for only a year or two, as long as you quit after that? Definitely no	n= 183 69.4 %+/- 6.7	67.7 %+/- 1.7

Jefferson County

Youth & Tobacco Data Summary 10th Grade Year 2000

	County %+/- Margin of Error	State %+/- Margin of Error
Would you ever use or wear something that has a tobacco company name or picture on it such as a lighter, t-shirt, hat, or sunglasses? Definitely no	n= 183 37.7 %+/- 7.0	39.7 %+/- 1.8
During last 12 months, did you buy or receive anything that has a tobacco name or picture on it? Yes	n= 184 22.3 %+/- 6.0	18.7 %+/- 1.4
During past 7 days, on how many days were you in the same room with someone who was smoking cigarettes? 3 or more days	n= 67 41.8 %+/- 11.8	28.7 %+/- 2.7
During past 7 days, on how many days did you ride in a car with someone who was smoking cigarettes? 3 or more days	n= 66 39.4 %+/- 11.8	24.6 %+/- 2.5
During this school year, did you practice ways to say 'NO' to tobacco in any of your classes (for example, by role playing)? Yes	n= 184 43.5 %+/- 7.2	27.5 %+/- 1.6
During past 30 days, have you seen or heard commercials on TV, the Internet, or on the radio about the dangers of cigarette smoking? More than once a week	n= 184 72.3 %+/- 6.5	77.6 %+/- 1.6
How wrong would most adults in your neighborhood think it was for kids your age to smoke cigarettes? Very wrong	n= 183 31.7 %+/- 6.7	39.4 %+/- 1.8
If you wanted to get some tobacco, how easy would it be for you to get some? Very hard or Sort of hard	n= 185 24.3 %+/- 6.2	27.8 %+/- 1.6
Has either of your parents (or guardians) discussed the dangers of tobacco use with you? Yes	n= 183 68.3 %+/- 6.7	65.8 %+/- 1.7
Does anyone who lives with you now smoke cigarettes? Yes	n= 67 43.3 %+/- 11.9	37.4 %+/- 2.9